

Cosmological Constraints on the Host Halos of GRBs

Wen Xu¹ and Li-Zhi Fang²

ABSTRACT

The recently observed bright optical transients(OT) of high redshift GRBs indicate that they are in a violent dynamical state. We think it is reasonable to assume that the GRBs form in the environment of gravitationally collapsed halos of the cosmic matter field, and we investigate the basic parameters of the halos which are favored to host GRBs. If the harboring coefficient f of GRBs per halo is weakly dependent on the mass of the halo, the redshift data of GRB OTs can yield significant constraints on the massive halos hosting GRBs. We show that, in the framework of popular cold dark matter (CDM) models, the GRB-favored environments are newly collapsed halos (i.e. their ages less than about 2×10^9 yr) with masses around $10^9 h^{-1} M_{\odot}$. In this scenario, low redshift GRBs, if they exist, could not have the same cosmic origin as the high redshift ones. To fit with the observed rate of GRBs, we conclude that each GRB halo can host probably no more than one GRB event on average. This result implies that GRBs may be related to the merging of the halos.

Subject headings: cosmology: theory - gamma rays: bursts - large-scale structure of universe

1. Introduction

The cosmological origin of γ -ray bursts (GRBs) is favored by many recent observations (Meegan et al. 1992; Costa et al. 1997; van Paradijs et al. 1997; Bond 1997; Fail et al. 1997; Metzger et al. 1997; Kulkarni et al. 1998; Djorgovski et al. 1998). The discovery of the bright optical transient (OT) associated with the GRB990123 (Odewahn et al. 1999a,b) at a redshift of $z=1.6$ (Kelson et al. 1999; Hjorth et al. 1999) added another cogent piece of evidence for the cosmological distances of GRBs. These results also lead to an “energy crisis” of about 3×10^{54} ergs if γ -ray emission is isotropic (Kulkarni et al. 1999). It is difficult to make a model of the energy source by known mechanisms of radiation, despite the many models of the GRB energy source which have been proposed (see the references in Hartmann 1996; Piran 1998). Nevertheless,

¹Department of Physics and Astronomy, Arizona State University, Tempe, AZ 85287

²Department of Physics, University of Arizona, Tucson, AZ 85721

the energy of GRBs indicates that they are related to some kind of violent process. Related to this idea, we note the galaxy host of GRB990123 is highly irregular (Bloom et al. 1999, Fruchter et al. 1999). This is evidence that the hosts of GRBs are in a violent dynamical state, such as gravitationally collapsed halos. Therefore, it would be reasonable to assume that GRBs form in collapsed halos during cosmic gravitational clustering. A question is then: What are the basic properties of these halos? Namely, in what environment (halos) are GRBs likely to form? We will try, in this *Letter*, to approach this question cosmologically.

The current situation of GRBs is in some ways similar to that of QSOs in the early 1970s: the mechanism of their radiation is uncertain, but they are believed to be cosmological objects. Past experience tells us that the cosmological aspects of QSOs can be studied even in the absence of knowledge of the radiation mechanism. For instance, the abundance, two-point correlation function, and redshift evolution of QSOs has been successfully investigated in the framework of large scale structure formation (Efsthathiou & Rees 1988, Nusser & Silk 1993, Blanchard, Bouchart & Klaff 1993, Bi & Fang, 1997). In this approach QSOs were assumed to be hosted by collapsed dark matter halos of the cosmic mass field, while the radiation and other hydrodynamic processes of QSOs were described by phenomenological parameters. Considering that the hydrodynamic processes are local, it is reasonable to assume that the hydrodynamic conditions may not be modulated by the density inhomogeneities on scale sizes larger than the size of the halo. In this case, the probability of a halo having a QSO should be the same for all halos (Fang & Jing 1998).

With the encouragement of the success of the cosmological approach to QSOs, we introduce a phenomenological parameter f to describe the average number of GRBs hosted by a collapsed halo. Consequently, the GRB birthrate at redshift z should be proportional to the birthrate of the considered collapsed halos. Thus, by comparing the cosmologically predicted halos with the observed redshifts of GRBs, one can find what halo environment is favored by GRBs, i.e. one can find the mass and age of the halos which are likely to host GRBs.

The current data on cosmological properties of GRBs are still poor. However, popular models of structure formation, such as the cold dark matter cosmogony, show that the redshift evolution of collapsed halos is a strong function of the parameters of the halos. In such a case, even a few redshift data are able to constrain the models. For instance, the data of three high redshift galaxy clusters were found to effectively rule out models with high mass density (Bahcall, Fan & Cen 1997). Therefore, one can expect that the few redshift observations of GRBs would also be able to provide useful constraints on the halos hosting GRBs. In the next two sections, using the data on the redshifts of GRB OTs and the rate of GRBs, we find the basic parameters of GRB-hosting halos in the framework of CDM cosmogony.

2. Redshift-dependence of the birthrate of GRBs

In the hierarchical structure formation models with Gaussian initial perturbation, the comoving number density of collapsed halos in the mass range M to $M + dM$ can be calculated with the Press & Schechter formalism (1974) as

$$n(M, z)dM = -\sqrt{\frac{2}{\pi}} \frac{\rho_0}{M^2} \frac{\delta_c}{\sigma(M, z)} \exp\left(\frac{-\delta_c^2}{2\sigma(M, z)^2}\right) \frac{d \ln \sigma(M, z)}{d \ln M} dM, \quad (1)$$

where $\delta_c \approx 1.69$ almost independent of cosmologies. $\sigma(M, z)$ is the linear theory *rms* mass density fluctuation in spheres of mass M at redshift z within a top-hat window of radius R , and is determined by the initial density spectrum $P(k)$ and normalization factor $\sigma_8 = \Delta(8 h^{-1} \text{Mpc}, 0)$.

We consider two popular representative CDM models - the standard CDM (SCDM) and the flat low-density CDM (LCDM). The parameters of the models (Hubble constant $h = H_0/(100 \text{ km s}^{-1} \text{ Mpc}^{-1})$, mass density Ω_0 , cosmological constant λ_0 and σ_8) are taken to be (0.5, 1, 0, 0.6) and (0.75, 0.3, 0.7, 1) for the SCDM and LCDM, respectively. These models provide a reasonable description of many observational properties of large scale structures of the universe.

Six high redshift galaxies have been found to host GRBs so far. Collapsed halos must be massive enough to be favored by the huge energy and violent activity of GRBs. Thus, we have to introduce the first parameter for the environment of GRBs: the mass scale M_{GRB} of GRB hosting halos. From eq.(1), the number density of the collapsed halos formed by the era z with mass greater than M_{GRB} is given by

$$N(> M_{GRB}, z) = \int_{M_{GRB}}^{\infty} n(M, z) dM. \quad (2)$$

Because $n(M, z)$ decreases rapidly with M , $N(> M_{GRB}, z)$ is actually dominated by halos with mass $M \simeq M_{GRB}$.

In the case of objects like galaxy clusters, there is a one-to-one correspondence between an object and a halo with mass M and radius r (Xu, Fang & Wu, 1998; Xu, Fang & Deng 1999). Obviously, the one-to-one identification is incorrect for the GRB as one massive collapsed halo may host many GRBs in its history. Moreover, the number of GRBs hosted by a halo of mass M may be different for different halos, because the formation of a GRB is not only determined by gravitational parameters of the halos, but also by hydrodynamical processes. To reduce this uncertainty, we consider that the hydrodynamical processes are local. This is, the relevant hydrodynamical conditions may not be modulated by the density inhomogeneities on scales much larger than the size of the halos considered. In this case, the average number of GRBs (f) hosted by a halo of mass M does not depend on the structures larger than the halos of eq.(2), and is also redshift-independent. Thus, the total number of GRBs hosted by halos of eq.(2) is $fN(> M_{GRB}, z)$.

The birthrate of halos of mass $M > M_{GRB}$ at z' is $dN(> M_{GRB}, z')/dt$. A GRB may not form at the same time as the birth of a collapsed halo of mass M . In either stellar models or

active galactic center models of GRBs, the formation of a GRB on average has to be later than the birth of its host by τ , which is the time scale of the evolution from the birth of a collapsed halo to the formation of GRBs in the halo. Obviously, τ is dependent on hydrodynamics, and different for different halos. However, similar to f , τ can be treated as a parameter if we are interested only in the mean rate of the GRB formations over a mass scale much larger than the GRB halos themselves. Thus, a halo of mass M_{GRB} , formed at redshift z' , will contribute on average f GRB(s) at epoch z , and

$$t(z) - t(z') = \tau. \quad (3)$$

The GRB birthrate at redshift z is then

$$\phi(z) = f \frac{dN(> M_{GRB}, z')}{dt} \frac{1 + z'}{1 + z}. \quad (4)$$

The last term comes from the time dilation between $z = z$ and $z = z'$.

Fig. 1 plots the birthrate $\phi(z)/f$ vs. z in models SCDM and LCDM for a given $M_{GRB} = 10^9 h^{-1} M_\odot$. For other values of M_{GRB} , the curves would basically be the same except for the normalization. The parameter τ is taken to be 0.1 to 5 Gyr. The curves in Fig.1 have similar features: sharply rising at redshift z_r and falling at z_f . Therefore, the nonzero range from z_r to z_f is weakly affected by the redshift dependence of f . The peak of the birth rate shifts to smaller redshift for larger τ . This is expected as the larger τ means the later formation of GRBs.

The observational data of GRB redshifts have been obtained in a couple of ways listed below with decreasing reliability: (1) Derived from absorption line features in the spectrum of optical transient. It includes the measurements of $z = 1.6$ for GRB990123 (Kelson et al. 1999; Hjorth et al. 1999) and $z = 0.835$ for GRB970508 (Metzger et al. 1997). (2) Derived from host galaxy, by assuming a physical association between GRB and its host galaxy. It includes the redshift measurements of $z = 0.835$ for GRB970508 (Bloom et al. 1998), $z = 0.966$ for GRB980703 (Djorgovski et al. 1998), $z = 1.096$ for GRB980613 (Djorgovski et al. 1999), $z = 3.418$ for GRB971214 (Kulkarni et al. 1998), and $1.3 \leq z \leq 2.5$ for GRB970228 (Fruchter et al. 1998). (3) Derived from a variety of objects other than OTs or host galaxies. Examples are GRB980329 at $z \sim 5$ (Fruchter 1999), GRB970828 at $z \sim 0.33$ (Yoshida et al. 1999) and GRB980425 at $z = 0.0085$ (Galama et al. 1998).

Our major conclusion is based only on the two OT redshifts. The OT redshift of GRB990123 ($z = 1.6$) is further considered as the mean of redshift distribution, because it is not far away from the middle value of the five available host galaxy redshifts. The redshift $z = 1.6$ is shown in Fig 1. This redshift is outside the birth rate curves with parameters of $\tau > 2 \times 10^9$ yr for SCDM, and $\tau > 3 \times 10^9$ yr for LCDM, regardless of M_{GRB} . However, if we take $z = 3.4$ of GRB971214 seriously, even $\tau > 1 \times 10^9$ yr curves can be excluded. These upper bounds of τ are significantly shorter than the Hubble time t_{Hubble} at redshift $z = 1.6$. ($t_{Hubble} = 3.1, 5.5$ Gyr, respectively for SCDM and LCDM). Therefore, GRBs are most likely to form in *newly* collapsed halos.

A generic feature of the birthrate of Fig. 1 is the deficiency of GRBs at low redshift. The

redshift range $z_r - z_f$ of each birthrate curve is generally rather narrow. z_r is larger than 0.1 for $\tau < 2 \times 10^9$ yr. Especially for the model LCDM, we have $z_r \gg 0.1$. Therefore, if the $z=0.0085$ supernova really corresponds to GRB980425 (Galama et al. 1998), the model LCDM will be in difficulty. Namely, GRB redshifts of $z = 1.6$ and 0.0085 can't be simultaneously contained within redshift range of $z_r - z_f$. The $z=0.0085$ supernova-related GRB probably does not belong to the same type of GRBs at high redshifts in our model. The redshift distribution also supports the speculation that the soft γ -ray repeaters in our galaxy are caused by neutron stars, and are different from GRBs with cosmic distances (Piran 1998).

Fig. 2 plots the birthrate $\phi(z)/f$ vs. z in models SCDM and LCDM for $\tau = 1$ and 2 Gyr, respectively for SCDM and LCDM. The parameter M_{GRB} is taken to be in the range $10^6 - 10^{10} h^{-1} M_\odot$. This figure also shows a sharply rising of the distribution at redshift z_r and a sharply falling at z_f . The peaks of the birth rate shift to smaller redshift for larger M_{GRB} , as the larger M_{GRB} formed later.

In order to fit the range of $z_r \simeq 1$ (GRB980703) and $z_f \simeq 3$ (GRB971214), the mass M_{GRB} of halos must be larger than $10^7 h^{-1} M_\odot$. Halos of $M_{GRB} \leq 10^7 h^{-1} M_\odot$ form too early, and it is difficult for them to simultaneously cover the $z \simeq 1$ and $z \simeq 3$ for any τ . On the other hand, $M_{GRB} > 10^{10} h^{-1} M_\odot$ halos form too late. Therefore, the most likely mass range of GRB halos is $M_{GRB} \simeq 10^7 - 10^{10} h^{-1} M_\odot$. This mass range is at the lower end of masses typical of today's galaxies. Therefore, these halos would be most readily identified as the progenitors of galaxies. This result is consistent with the observed hosts, as the mass of GRB hosts is on the order of the mass of halos that can form galaxies by merging. It doesn't mean, however, that the hosts of GRBs *must* be galaxies, because some massive halos may not be involved in galaxy formation. For these halos GRBs are not associated with galaxies.

3. Number counts of GRBs

We now consider the reasonableness of the environment parameters with the observed rate of GRB. From the birthrate eq.(4), the all-sky, observable number per year, of gamma ray burst events at the epoch of redshift z is

$$N_{GRB}(z)dz = \phi(z)(1+z)dV, \quad (5)$$

where $dV(z)$ is the volume element in redshift space. The term $(1+z)$ comes from the time dilation between $z = 0$ and $z = z$. Thus, the total number of GRBs per year is

$$N_{GRB} \equiv \int N_{GRB}(z)dz = f \cdot N_{halo}, \quad (6)$$

where

$$N_{halo} \equiv \int \frac{dN(> M_{GRB}, z')}{dt} (1+z') \frac{dV}{dz} dz. \quad (7)$$

The observed GRBs per unit time is then

$$N_{obs} = s N_{GRB} \frac{\delta\Omega}{4\pi} = s f \frac{\delta\Omega}{4\pi} N_{halo}, \quad (8)$$

where s is a factor less than 1 to describe observational efficiency of GRBs, $\delta\Omega/4\pi$ is the correction of angular scale if the radiation is beamed.

The all-sky observed rate of GRBs is approximately one per day, i.e. $N_{obs} \simeq 400\text{yr}^{-1}$. Thus, one can calculate the observation-theory ratio N_{obs}/N_{halo} , which is equal to $s f \delta\Omega/4\pi$. Fig. 3 shows how N_{obs}/N_{halo} depends on M_{GRB} and τ . It is interesting to see that for all the considered parameter ranges of M_{GRB} and τ , N_{obs}/N_{halo} is less than, and even far less than one.

Because $s \leq 1$ and $\delta\Omega/4\pi \leq 1$, we have a lower bound to f given by

$$f > \frac{N_{obs}}{N_{halo}}. \quad (9)$$

Since $M_{GRB} \geq 10^7 h^{-1} M_\odot$ (§2) and $\tau \leq 2$ Gyr (SCDM) and 3 Gyr (LCDM), we have $f > 3 \times 10^{-5}$ (LCDM) and 3×10^{-6} (SCDM).

There is some speculation that GRBs may constitute a unique homogeneous population of sources which has few selection effect (Piran 1998). If we further assume that s is not much less than 1, then $f \simeq N_{obs}/(N_{halo} \delta\Omega/4\pi)$. Although our guesses about the beaming of the γ -ray emission varying from isotropic explosion $\delta\Omega/4\pi = 1$ (e.g. Iwamoto et al. 1998) to highly beamed radiation $\delta\Omega = 0.1 - 0.001$ (e.g. Hartmann 1996), f is generally not larger than 1 for the majority of value of M_{GRB} and τ shown in Fig. 3. Our model predicts one new born massive halo can contribute no more than one GRB. A possible interpretation of this result is that the GRB phenomenon is related to halo merging. In the hierarchical structure formation scenario, massive halos formed from merging of less massive halos, i.e. each merging of massive halos will produce a new halo. Thus, each GRB halo will undergo no more than once of merging of itself. Imaging of GRB990123 shows that the GRB is apparently within an ongoing merger (Fruchter et al. 1999).

Moreover, if most GRB-hosted halos merged into galaxies, each galaxy with mass $10^{12} h^{-1} M_\odot$ formed, on average, from 10^5 merging of $10^7 h^{-1} M_\odot$ halos; or 10^2 merging of $10^{10} h^{-1} M_\odot$ halos. In other words, in the entire evolutionary history of galaxy formation, a galaxy on average underwent about $10^2 - 10^5$ GRB-hosted halo mergings. On the other hand, an estimation of the mean GRB number per galaxy in their evolutionary history found $\sim 10^4/\delta\Omega$ (Hartmann 1996), which is consistent with our cosmological estimation.

4. Conclusions

The redshift data of GRB OTs yield significant constraints on the massive halos hosting GRBs if the popular models of CDM cosmogony are employed. We show that these data can

be fitted by standard CDM models. Our model predicts GRBs probably to be new members of the family of high redshift objects which trace the cosmic large scale structures. We show that the halos forming GRBs are newly collapsed ($\tau < 2 \times 10^9$ yr) with masses scales around $10^9 h^{-1} M_{\odot}$. If the current record of GRBs is free of observational selection effects, the observed rate of GRBs requires that each GRB host halo will host no more than one GRB event on average. This strongly implies a possible relation between halo mergings and GRBs.

These conclusions are based on our assumption of a constant harboring coefficient f . More generally, f may depend on the mass M of the collapsed halo. For instance, the capacity of harboring GRBs can be made proportional to M (i.e. a linear relation: $f = \text{constant} \cdot M$). Since the abundance of halos $N(z)$ is a strong function of redshift around z_r and z_f , any weak M -dependent relation of f will not change the conclusion about z_r and z_f . In this case, the constraints to M and τ are still valid.

Using the parameters of the possible GRB halos, we predict the redshift distribution of GRB OTs, as shown in Figs. 1 and 2. This redshift distribution is not sensitive to how f depends on M . Therefore, our results will be further tested in near future when enough redshift data are accumulated to give us a reliable redshift distribution of GRBs.

We would like to thank an anonymous referee for a report which improved the presentation of the paper. WX thanks H. Yan, T. Lu, R. Windhorst, D. Burstein, R. Clawson for valuable discussions.

REFERENCES

- Bahcall, N.A., Fan, X. & Cen, R.Y. 1997, ApJ, 485, L53.
- Bi, H.G. & Fang, L.Z. 1997, A&A, 325, 433.
- Blanchard, A., Buchartet, T., Klaff, R. 1993, A&A, 267, 1.
- Bloom, J. S., Djorgovski, S. G., Kulkarni, S. R., & Frail, D. A. 1998, ApJ, 507, L25.
- Bloom, J. S., Odewahn, S. C., Djorgovski, S. G., et al. 1999, astro-ph/9902182.
- Bond, H. E., 1997, IAU circ. 6665.
- Cohen, E., & Piran, T. 1995, ApJ, 444, 25.
- Costa, E., et al., 1997, Nature, 387, 783.
- Djorgovski, S. G., et al., 1998, CGN notice 139.
- Djorgovski, S. G., et al., 1999, GCN Circ. 189.
- Efstathiou, G. & Rees, M. 1988, MNRAS, 230, 5p.
- Eke, V.R., Cole, S., Frenk, C.S. & Navarro, J.F. 1996, MNRAS, 281, 703.

- Fang, L.Z. & Jing, Y.P. 1998, ApJ, 502, L95.
- Frail, D. A., et al., 1997, Nature, 389, 261.
- Fruchter, A.S. 1999, ApJ, 512, L1.
- Fruchter, A.S. et al. 1998, ApJ, in press, (astro-ph/9807295).
- Fruchter, A.S. et al. 1999, ApJ, in press, (astro-ph/9902236).
- Hartmann, D. H. 1996, Astron. Astrophys. Suppl. Ser. 120, 31.
- Hjorth, J. et al., 1999, GCN Circular 219.
- Iwamoto, K. et al. 1998, Nature, 395, 672.
- Galama, T. J. et al. 1998, Nature, 395, 670.
- Kelson, D. D., Illingworth, G. I., Franx, M., Magee, D. & van Dokkum, P. G., 1999, IAU Circular 7096.
- Kulkarni, S., et al., 1998, Nature, 393, 35.
- Meegan, C. A., et al., 1992, Nature, 355, 143.
- Metzger, M. R., Djorgovski, S. G., Kulkarni, S. R. et al. 1997, Nature 387, 878.
- Nusser, A. & Silk, J. 1993, ApJ, 411, L1.
- Odewahn, S. C., Bloom, J. S., Kulkarni, S. R. 1999a, G.C.N. 201.
- Odewahn, S. C., Bloom, J. S., Kulkarni, S. R. 1999b , I.A.U.C. 7094.
- Piran, T. 1998, to appear in Physics Reports, (astro-ph/9810256.)
- Press, W.H. & Schechter, P. 1974, ApJ, 187, 425.
- van Paradijs, J., et al., 1997, Nature, 386, 686.
- Xu, W., Fang, L.Z., & Wu, X.P. 1998, ApJ, 508, 472.
- Xu, W., Fang, L.Z., & Deng, Z.G. 1999 ApJ, in press.
- Yoshida, A. et al. 1999, in Proceedings of the Gamma-ray Bursts in the Afterglow Era Workshop, Rome, Nov 1998, A&AS, in press.

Fig. 1.— The redshift evolution of the birth rate of GRBs. The ratio $\phi(z)/f$ is shown so that the curves do not depend on the choice of f . The solid and dashed lines are for LCDM and SCDM, respectively. The parameter τ is taken to be 5, 3, 2, 1, 0.1 Gyr (LCDM) and 5, 3, 1, 0.1 Gyr (SCDM) from left to right. The parameter M_{GRB} is fixed at $10^9 h^{-1} M_{\odot}$. The redshift of GRB990123 is indicated as a landmark of z . The model curves with the best choice by the redshift data of GRBs are highlighted as thick lines.

Fig. 2.— The redshift evolution of the birth rate of GRBs. The solid and dashed lines are for LCDM and SCDM, respectively. The mass of GRB parent halos are taken to be $10^6, 10^7, 10^8, 10^9, 10^{10} h^{-1} M_{\odot}$ from top to bottom curves, respectively. τ is taken to be 2×10^9 years for LCDM and 1×10^9 years for SCDM. The redshift of GRB990123 is indicated as a landmark of z .

Fig. 3.— The ratio N_{obs}/N_{halo} as a function of parameters τ and M_{GRB} . The solid lines are for LCDM and dashed lines for SCDM. The parameter τ is equal to 5, 3, 2, 1, 0.1 Gyr from top to bottom. The highlighted two lines are the same as the two in Fig.1.





